



# SOLVABILITY OF THE THIRD-ORDER KORTEWEG-DE VRIES (KDV) EQUATION BY VARIATIONAL ITERATION AND NEW ITERATIVE METHODS

*\*Usman, M.A., Shittu, M.T., Solanke, O.O and Hammed F.A.*

*Department of Mathematical Sciences  
Olabisi Onabanjo University, Ago –Iwoye, Nigeria*

*\*Corresponding Email: usman.mustapha@oouagoiwoye.edu.ng*

*Manuscript Received: 15/06/2020    Accepted: 11/08/2020    Published: December, 2020*

## ABSTRACT

This paper examined the approximate solution of the third-order Korteweg-de Vries (KdV) equations is obtained by the Variational Iteration Method (VIM) developed by Ji-Huan He and the New Iterative Method (NIM) developed by Daftardar Gejji and Jafari. These methods provide the solution in the form of a convergent series which illustrate the ability and the effectiveness of the methods, some examples were provided. The results showed that the methods are very simple, effective, powerful and can easily be applied to other linear and nonlinear PDEs.

**Keywords:** *Variational iteration Method, New Iterative Method, Korteweg-de Vries equations, Lagrange multiplier.*

**INTRODUCTION**

The Korteweg-de Vries equation (KdV) was derived by D.J. Korteweg and G. de Vries in 1895. The KdV equation is used to study the unusual water waves that occur in shallow, narrow channels such as canals (Olusola & Oluwole, 2012). It also occurs in many physical applications such as non-linear plasma waves which exhibit certain dissipative effects, propagation of waves and propagation of bores in shallow water waves. (Muhammad *et al.*, 2019)

The third order Korteweg-de Vries (KdV) equation is of the form:

$$u_t(x,t) - auu_x(x,t) + bu_{xxx}(x,t) = 0 \quad \dots\dots(1)$$

Subject to the initial conditions

$$u(x,0)=f(x) \quad \dots\dots\dots(2)$$

where a and b are constants,  $u_x$  and  $u_t$  are partial derivatives of function u with respect to space x and t respectively and the non-linear term  $uu_x$  tends to localize the wave, whereas the wave was spread out by dispersion. The formulation of solitons that have a single humped wave was define by delicate balance between  $uu_x$  and  $u_{xxx}$ . the displacement which described how waves evolve under the competing but comparable effects of weak nonlinearity and weak dispersion is denoted by  $u(x,t)$  [Olufemi *et al.*, 2019]

Over the years, Scientists have made several attempts to obtain the analytical and numerical solutions of KdV equation, Some of the methods used are Homotopy Perturbation Method (Eljaily & Tarig, 2015), Elzaki Transform Method (Shraddha & Mihir, 2014), Adomian Polynomial and Elzaki Transform Method (Olufemi *et al.*, 2019), New Cubic B-Spline Method (Muhammad *et al.*, 2019), Hyperbola Function Method (Luwai, 2016), Finite Difference Method and Adomian Decomposition Method (Olusola & Oluwole, 2012), Multiquadric Quasi-Interpolation Method (Xiao *et al.*, 2011; Maryam & Azim, 2015), Meshfree Method (Siraj *et al.*, 2008), Collocation and Radial Basis Function Method (Dehghan and Shokri, 2007), Method of Lines (Schuesser, 1994), Small Time Solution (Kutluay *et al.*, 2000), Finite Difference Method (Bahadir, 2005), Homotopy Analysis Method (Vincent *et al.*, 2013), Variational Iteration Method, Homotopy Perturbation and Homotopy Analysis Method (Foad *et al.*, 2010) and many others.

In this paper, the solution of third order Korteweg-de Vries (KdV) equations are obtained by Variational Iteration Method (VIM) and New Iterative Method (NIM). These methods give the solutions as an approximate analytical solution in series form and it yields exact solution with few iterations.

**VARIATIONAL ITERATION METHOD (VIM)**

To clarify the basic ideas of VIM, we consider the following differential equation:

$$Lu+Nu=g(t) \quad \dots\dots\dots(3)$$

Where, L and N are linear and nonlinear operators respectively, and g(t) is the source inhomogeneous term.

The variational iteration method presents a correction functional as follows:

$$u_{n+1}(t) = u_n(t) + \int_a^b \lambda(\varepsilon)(Lu_n(\varepsilon) + N\tilde{u}_n(\varepsilon) - g(\varepsilon))d\varepsilon \quad \dots\dots(4)$$

where  $\lambda$  is a general Lagrange multiplier, which can be identified optimally via the variational theory, and  $\tilde{u}_n$  is a restricted variation which means  $\tilde{u}_n = 0$ .

It is obvious now that the main steps of the variational iteration method require first the determination of the Lagrange multiplier  $\lambda(\varepsilon)$  that will be identified optimally. Integration by parts is usually used for the determination of the Lagrange multiplier  $\lambda(\varepsilon)$ . In other words, carrying out the integration as follows can yield:

$$\int \lambda(\varepsilon) u'_n(\varepsilon) d\varepsilon = \lambda(\varepsilon) u_n(\varepsilon) - \int \lambda'(\varepsilon) u_n(\varepsilon) d\varepsilon \quad \dots\dots 5$$

Having determined the Lagrange multiplier  $\lambda(\varepsilon)$ , the successive approximations  $u_{n+1}, n \geq 0$ , of the solution u will be readily obtained upon using any selective function  $u_0$ .

However, for fast convergence, the function  $u_0(x,t)$  should be selected by using the initial conditions as follows:

$$u_0(x,t)=u(x,0) \quad \text{for first order}$$

Consequently, the solution

$$u = \lim_{n \rightarrow \infty} u_n \quad \dots\dots\dots 6$$

In other words, equation (5) will give several approximations, and therefore the exact solution is obtained as the limit of the resulting successive approximations (Batiha, 2009) (Wazwaz, 2009).

**NEW ITERATIVE METHOD (NIM)**

To illustrate the idea of the NIM, we consider the following general functional equation:

$$u=f+N(u) \quad \dots\dots\dots(7)$$

where N is a nonlinear operator and f is a given function. We can find the solution of equation (7) having the series of the form

$$u = \sum_{i=0}^{\infty} u_i \quad \dots\dots\dots(8)$$

The nonlinear operator N can be decomposed as:

$$N(\sum_{i=0}^{\infty} u_i) = N(u_0) + \sum_{i=1}^{\infty} \left\{ N\left(\sum_{j=0}^i u_j\right) - N\left(\sum_{j=0}^{i-1} u_j\right) \right\} \quad \dots\dots 9$$

Substituting equations (8) and (9) into equation (7) gives

$$\sum_{i=0}^{\infty} u_i = f + N(u_0) + \sum_{i=1}^{\infty} \left\{ N\left(\sum_{j=0}^i u_j\right) - N\left(\sum_{j=0}^{i-1} u_j\right) \right\} \dots\dots 10$$

$$\sum_{i=0}^m u_i = f + N\left(\sum_{j=0}^m u_j\right) \dots\dots\dots 12$$

We define the recurrence relation of equation in the following way:

$$\begin{aligned} u_0 &= f \\ u_1 &= N(u_0) \\ u_2 &= N(u_0 + u_1) - N(u_0) \dots\dots\dots (11) \\ u_3 &= N(u_0 + u_1 + u_2) - N(u_0 + u_1) \\ u_{n+1} &= N(u_0 + u_1 + \dots + u_n) - N(u_0 + u_1 + \dots + u_{(n-1)}); \quad n=1, 2, 3 \end{aligned}$$

Then  
 $u_1 + \dots + u_{m+1} = N(u_0 + u_1 + \dots + u_m); \quad m = 1, 2, 3$   
 and

The m-term approximate solution of (7) is given by  
 $u = u_0 + u_1 + u_2 + \dots + u_{m-1}$   
 (Bhaleka S. & Daftardar-Gejji V.)

**NUMERICAL EXAMPLES**

Example 4.1

Consider the following KDV equation

$$u_t - 6uu_x + u_{xxx} = 0 \dots\dots\dots 13$$

With the initial condition

$$u(x, 0) = 6x$$

The exact solution is given by  $u(x, t) = \frac{6x}{1 - 36t}$

**Method 1: VIM**

The correction functional for example 4.1 is

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \lambda(c) \left[ \frac{\partial u_n(x, \varepsilon)}{\partial \varepsilon} - 6\tilde{u}_n(x, c) \frac{\partial \tilde{u}_n(x, \varepsilon)}{\partial x} + \frac{\partial^3 \tilde{u}_n(x, \varepsilon)}{\partial x^3} \right] dc \dots\dots\dots 14$$

Taking variations on both sides gives

$$\delta u_{n+1}(x, t) = \delta u_n(x, t) + \int_0^t \delta \lambda(c) \left[ \frac{\partial u_n(x, \varepsilon)}{\partial \varepsilon} - 6\tilde{u}_n(x, c) \frac{\partial \tilde{u}_n(x, \varepsilon)}{\partial x} + \frac{\partial^3 \tilde{u}_n(x, \varepsilon)}{\partial x^3} \right] dc$$

$\tilde{u}_n$  is a restrictive vibration, therefore  $\delta \tilde{u}_n = 0$

$$\delta u_{n+1}(x, t) = \delta u_n(x, t) + \int_0^t \delta \lambda(c) \left[ \frac{\partial u_n(x, \varepsilon)}{\partial \varepsilon} \right] dc$$

*Simplify by integration by parts*

$$\delta u_{n+1}(x, t) = \delta u_n(x, t) + \delta \lambda(c) u_n(x, c) - \int_0^t \lambda'(c) \delta u_n(x, c) dc$$

Making the correction functional stationary

$$\delta u_{n+1}(x, t) = 0$$

We have

$$0 = \delta u_n(x, t) + \delta \lambda(c) u_n(x, c) - \int_0^t \lambda'(c) \delta u_n(x, c) dc$$

The stationary points are  $1 + \lambda = 0$  and  $\lambda' = 0$

Solving this, we have  $\lambda = -1$

Hence, the iterative formula becomes

$$u_{n+1}(x, t) = u_n(x, t) - \int_0^t \left[ \frac{\partial u_n(x, \varepsilon)}{\partial \varepsilon} - 6u_n(x, c) \frac{\partial u_n(x, \varepsilon)}{\partial x} + \frac{\partial^3 u_n(x, \varepsilon)}{\partial x^3} \right] dc \dots\dots\dots 15$$

From the initial condition, we have

$$u_0 = 6x$$

Equation (15) at  $n = 0$  becomes

$$u_1(x, t) = u_0(x, t) - \int_0^t \left[ \frac{\partial u_0(x, \varepsilon)}{\partial \varepsilon} - 6u_0(x, c) \frac{\partial u_0(x, \varepsilon)}{\partial x} + \frac{\partial^3 u_0(x, \varepsilon)}{\partial x^3} \right] dc$$

$$u_1(x, t) = 6x + 6^3xt = 6x(1 + 36t)$$

When n = 1

$$u_2(x, t) = u_1(x, t) - \int_0^t \left[ \frac{\partial u_1(x, \varepsilon)}{\partial \varepsilon} - 6u_1(x, c) \frac{\partial u_1(x, \varepsilon)}{\partial x} + \frac{\partial^3 u_1(x, \varepsilon)}{\partial x^3} \right] dc$$

$$u_2(x, t) = 6x + 6^3t + 6^5xt^2 = 6x(1 + 36t + (36t)^2)$$

Consequently, the successive approximations are obtained

$$u_1(x, t) = 6x(1+36t)$$

$$u_2(x, t) = 6x(1+36t+(36t)^2)$$

$$u_3(x, t) = 6x(1+36t+(36t)^2+(36t)^3)$$

Hence, the closed form and exact solution is given as

$$u(x, t) = 6x(1+36t+(36t)^2+(36t)^3+(36t)^4+\dots) \dots 16$$

$$u(x, t) = \frac{6x}{1 - 36t} \dots 17$$

Method 2: NIM

Considering Example 4.1

Using NIM, the equation (13) is equivalent to the following integral equation

$$u = f + \int_0^t (6uu_x - u_{xxx}) dt \dots 18$$

From the initial condition, we have

$$u_0 = 6x$$

$$N(u_k) = \int_0^t (6u_k u_{kx} - u_{kxxx}) dt$$

When k = 0,

$$N(u_0) = u_1 = \int_0^t (6u_0 u_{0x} - u_{0xxx}) dt$$

$$= \int_0^t (6(6x)(6) - 0) dt = \int_0^t 6^3 x dt = 6^3 xt$$

$$u_1 = 6^3 xt$$

$$u_2 = N(u_0 + u_1) - N(u_0)$$

$$u_2 = 6^3 xt + 6^5 xt^2 - 6^3 xt = 6^5 xt^2$$

$$u_3 = 6^3 xt + 6^5 xt^2 + 6^7 xt^3 - 6^3 xt - 6^5 xt^2 = 6^7 xt^3$$

The successive approximations are:

$$u_0 = 6x$$

$$u_1 = 6^3 xt$$

$$u_2 = 6^5 xt^2$$

$$u_3 = 6^7 xt^3$$

Thus the series solution of example 4.1 is given by

$$u(x, t) = \sum_{n=1}^{\infty} u_n = 6x + 6^3 xt + 6^5 xt^2 + 6^7 xt^3$$

$$u(x, t) = 6x + 6^3 xt + 6^5 xt^2 + 6^7 xt^3$$

$$u(x, t) = 6x(1+36t+(36t)^2+(36t)^3+(36t)^4+\dots)$$

$$u(x, t) = \frac{6x}{1 - 36t} \dots 19$$




Figure 1: The exact solution of example 4.1 ( $u(x,t)=6x/(1-36t)$ )

Example 4.2

Consider the following KDV equation

$$u_t + 6uu_x + u_{xxx} = 0 \dots 20$$

With the initial condition

$$u(x, 0) = x$$

The exact solution is given by  $u(x, t) = \frac{x}{1 + 6t}$

Method 1: VIM

The correction functional for example 4.2 is

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \lambda(c) \left[ \frac{\partial u_n(x, \varepsilon)}{\partial \varepsilon} + 6\tilde{u}_n(x, c) \frac{\partial \tilde{u}_n(x, \varepsilon)}{\partial x} + \frac{\partial^3 \tilde{u}_n(x, \varepsilon)}{\partial x^3} \right] dc \dots 21$$

Taking variations on both sides gives

$$\delta u_{n+1}(x, t) = \delta u_n(x, t) + \int_0^t \delta \lambda(c) \left[ \frac{\partial u_n(x, \varepsilon)}{\partial \varepsilon} + 6\tilde{u}_n(x, c) \frac{\partial \tilde{u}_n(x, \varepsilon)}{\partial x} + \frac{\partial^3 \tilde{u}_n(x, \varepsilon)}{\partial x^3} \right] dc$$

$\tilde{u}_n$  is a restrictive vibration, therefore  $\delta \tilde{u}_n = 0$

$$\delta u_{n+1}(x, t) = \delta u_n(x, t) + \int_0^t \delta \lambda(c) \left[ \frac{\partial u_n(x, \varepsilon)}{\partial \varepsilon} \right] dc$$

From previous calculation,  $\lambda = -1$

Hence, the iterative formula becomes

$$u_{n+1}(x, t) = u_n(x, t) - \int_0^t \left[ \frac{\partial u_n(x, \varepsilon)}{\partial \varepsilon} + 6u_n(x, c) \frac{\partial u_n(x, \varepsilon)}{\partial x} + \frac{\partial^3 u_n(x, \varepsilon)}{\partial x^3} \right] dc$$

From the initial condition, we have

$$u_0 = x$$

When n = 0

$$u_1(x,t) = u_0(x,t) - \int_0^t \left[ \frac{\partial u_0(x,\varepsilon)}{\partial \varepsilon} + 6u_0(x,\varepsilon) \frac{\partial u_0(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_0(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

$$u_1(x,t) = x - \int_0^t 6x d\varepsilon = x - 6xt$$

When n = 1

$$u_2(x,t) = u_1(x,t) - \int_0^t \left[ \frac{\partial u_1(x,\varepsilon)}{\partial \varepsilon} + 6u_1(x,\varepsilon) \frac{\partial u_1(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_1(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

$$u_2(x,t) = x - 6xt + 6^2 xt^2$$

$$u_3(x,t) = x - 6xt + 6^2 xt^2 - 6^3 xt^3$$

Consequently, the successive approximations are obtained

$$u_1(x,t) = x(1-6t)$$

$$u_2(x,t) = x(1-6t+(6t)^2)$$

$$u_3(x,t) = x(1-6t+(6t)^2-(6t)^3)$$

Hence, the closed form and exact solution is given as

$$u(x,t) = x(1-6t+(6t)^2-(6t)^3+(6t)^4+\dots)$$

$$u(x,t) = \frac{x}{1+6t} \dots\dots\dots 22$$

Method 2: NIM

Considering Example 4.2

Using NIM, the equation is equivalent to the following integral equation

$$u = f + \int_0^t (-6uu_x - u_{xxx}) dt \dots\dots\dots 23$$

From the initial condition, we have

$$u_0 = x$$

$$N(u_k) = \int_0^t (-6u_k u_{k,x} - u_{k,xxx}) dt$$

When k = 0,

$$N(u_0) = u_1 = \int_0^t (-6u_0 u_{0,x} - u_{0,xxx}) dt$$

$$= \int_0^t (-6x) dt = -6xt$$

$$u_1 = -6xt$$

$$u_2 = N(u_0 + u_1) - N(u_0)$$

$$u_2 = -6xt + 6^2 xt^2 + 6xt = 6^2 xt^2$$

$$u_3 = -6^3 xt^3$$

The successive approximations are:

$$u_0 = x$$

$$u_1 = -6xt$$

$$u_2 = 6^2 xt^2$$

$$u_3 = -6^3 xt^3$$

Thus, the series solution of example 4.2 is given by

$$u(x,t) = \sum_{n=1}^{\infty} u_n = x - 6xt + 6^2 xt^2 - 6^3 xt^3$$

$$u(x,t) = x - 6xt + 6^2 xt^2 - 6^3 xt^3$$

$$u(x,t) = x(1-6t+(6t)^2-(6t)^3+(6t)^4+\dots)$$

$$u(x,t) = \frac{x}{1+6t} \dots\dots\dots 24$$




Figure 2: The exact solution of example 4.2 ( $u(x,t)=x/1+6t$ )

Example 4.3

Consider the following KDV equation

$$u_t - 6uu_x + u_{xxx} = 0 \dots\dots\dots (25)$$

With the initial condition

$$u(x,0) = \frac{2}{x^2}$$

The exact solution is given by  $u(x,t) = \frac{2}{x^2}$

Method 1: VIM

The correction functional for example 4.3 is

$$u_{n+1}(x,t) = u_n(x,t) + \int_0^t \lambda(c) \left[ \frac{\partial u_n(x,\varepsilon)}{\partial \varepsilon} - 6u_n(x,\varepsilon) \frac{\partial u_n(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_n(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

Taking variations on both sides gives

$$\delta u_{n+1}(x,t) = \delta u_n(x,t) + \int_0^t \delta \lambda(c) \left[ \frac{\partial u_n(x,\varepsilon)}{\partial \varepsilon} - 6u_n(x,\varepsilon) \frac{\partial u_n(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_n(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

$\delta \bar{u}_n$  is a restrictive vibration, therefore  $\delta \bar{u}_n = 0$

$$\delta u_{n+1}(x,t) = \delta u_n(x,t) + \int_0^t \delta \lambda(c) \left[ \frac{\partial u_n(x,\varepsilon)}{\partial \varepsilon} \right] d\varepsilon$$

From previous calculation,  $\lambda = -1$

Hence, the iterative formula becomes

$$u_{n+1}(x,t) = u_n(x,t) - \int_0^t \left[ \frac{\partial u_n(x,\varepsilon)}{\partial \varepsilon} - 6u_n(x,\varepsilon) \frac{\partial u_n(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_n(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

From the initial condition, we have

$$u_0 = \frac{2}{x^2}$$

When n = 0

$$u_1(x,t) = u_0(x,t) - \int_0^t \left[ \frac{\partial u_0(x,\varepsilon)}{\partial \varepsilon} - 6u_0(x,\varepsilon) \frac{\partial u_0(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_0(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

$$u_1(x,t) = \frac{2}{x^2} - \int_0^t (48x^{-5} - 48x^{-5}) d\varepsilon = \frac{2}{x^2}$$

When  $n = 1$

$$u_2(x,t) = u_1(x,t) - \int_0^t \left[ \frac{\partial u_1(x,\varepsilon)}{\partial \varepsilon} + 6u_1(x,\varepsilon) \frac{\partial u_1(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_1(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

$$u_2(x,t) = \frac{2}{x^2} - \int_0^t (18x^{-5} - 18x^{-5}) d\varepsilon = \frac{2}{x^2}$$

$$u_3(x,t) = \frac{2}{x^2} - \int_0^t (18x^{-5} - 18x^{-5}) d\varepsilon = \frac{2}{x^2}$$

Consequently, the successive approximations are obtained

$$u_1(x,t) = \frac{2}{x^2}$$

$$u_2(x,t) = \frac{2}{x^2}$$

$$u_3(x,t) = \frac{2}{x^2}$$

Hence, the closed form and exact solution is given as  $u(x,t) = \frac{2}{x^2}$  .....26

**Method 2: NIM**

Example 4.3  
Using NIM, the equation is equivalent to the following integral equation

$$u = f + \int_0^t (6uu_x - u_{xxx}) dt$$

.....27

From the initial condition, we have

$$u_0 = \frac{2}{x^2}$$

$$N(u_k) = \int_0^t (6u_k u_{kx} - u_{kxxx}) dt$$

When  $k = 0$ ,

$$N(u_0) = u_1 = \int_0^t (6u_0 u_{0x} - u_{0xxx}) dt$$

$$= \int_0^t (-18x^{-5} + 18x^{-5}) dt = 0$$

$$u_1 = 0$$

$$u_2 = N(u_0 + u_1) - N(u_0)$$

$$u_2 = \int_0^t (-18x^{-5} + 18x^{-5}) dt = 0$$

$u_3 = 0$   
The successive approximations are:

$$u_0 = \frac{2}{x^2}$$

$$u_1 = 0$$

$$u_2 = 0$$

$$u_3 = 0$$

Thus, the series solution of example 4.3 is given by

$$u(x,t) = \sum_{n=1}^{\infty} u_n = \frac{2}{x^2} + 0 + 0 + 0 + \dots$$

$$u(x,t) = \frac{2}{x^2}$$

.....28




Figure 3: The exact solution of example 4.3 ( $u(x,t)=2/x^2$ )

Example 4.4  
Consider the following KDV equation  $u_t - 6uu_x + u_{xxx} = 0$  .....(29)

With the initial condition

$$u(x,0) = \frac{2}{(x-3)^2}$$

The exact solution is given by  $u(x,t) = \frac{2}{(x-3)^2}$

**Method 1: VIM**

The correction functional for example 4.4 is

$$u_{n+1}(x,t) = u_n(x,t) + \int_0^t \lambda(\varepsilon) \left[ \frac{\partial u_n(x,\varepsilon)}{\partial \varepsilon} - 6u_n(x,\varepsilon) \frac{\partial u_n(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_n(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

Taking variations on both sides gives

$$\delta u_{n+1}(x,t) = \delta u_n(x,t) + \int_0^t \delta \lambda(\varepsilon) \left[ \frac{\partial u_n(x,\varepsilon)}{\partial \varepsilon} - 6u_n(x,\varepsilon) \frac{\partial u_n(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_n(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

$\bar{u}_n$  is a restrictive vibration, therefore  $\delta \bar{u}_n = 0$

$$\delta u_{n+1}(x,t) = \delta u_n(x,t) + \int_0^t \delta \lambda(\varepsilon) \left[ \frac{\partial u_n(x,\varepsilon)}{\partial \varepsilon} \right] d\varepsilon$$

From previous calculation,  $\lambda = -1$   
Hence, the iterative formula becomes

$$u_{n+1}(x,t) = u_n(x,t) - \int_0^t \left[ \frac{\partial u_n(x,\varepsilon)}{\partial \varepsilon} - 6u_n(x,\varepsilon) \frac{\partial u_n(x,\varepsilon)}{\partial x} + \frac{\partial^3 u_n(x,\varepsilon)}{\partial x^3} \right] d\varepsilon$$

From the initial condition, we have

$$u_0 = \frac{2}{(x-3)^2}$$

$$u_1(x,t) = \frac{2}{(x-3)^2} - \int_0^t (18(x-3)^{-5} - 18(x-3)^{-5}) d\varepsilon = \frac{2}{(x-3)^2}$$

$$u_2(x,t) = \frac{2}{(x-3)^2} - \int_0^t (18(x-3)^{-5} - 18(x-3)^{-5}) d\varepsilon = \frac{2}{(x-3)^2}$$

$$u_3(x,t) = \frac{2}{(x-3)^2} - \int_0^t (18(x-3)^{-5} - 18(x-3)^{-5}) d\varepsilon = \frac{2}{(x-3)^2}$$

Thus the series solution of example 4.4 is given by

$$u_3(x,t) = \frac{2}{(x-3)^2} - \int_0^t (18(x-3)^{-5} - 18(x-3)^{-5}) dt = \frac{2}{(x-3)^2}$$

$$u(x,t) = \sum_{n=1}^{\infty} u_n = \frac{2}{(x-3)^2} + 0 + 0 + 0 + \dots$$

$$u(x,t) = \frac{2}{(x-3)^2}$$

Consequently, the successive approximations are

$$u_0(x,t) = \frac{2}{(x-3)^2}$$

$$u_1(x,t) = \frac{2}{(x-3)^2}$$

$$u_2(x,t) = \frac{2}{(x-3)^2}$$

$$u_3(x,t) = \frac{2}{(x-3)^2}$$

Hence, the closed form and exact solution is given as

$$u(x,t) = \frac{2}{(x-3)^2} \dots\dots\dots 30$$

**Method 2: NIM**

Example 4.4

Using NIM, the equation is equivalent to the following integral equation

$$u = f + \int_0^t (6uu_x - u_{xxx}) dt \dots\dots\dots 31$$

From the initial condition, we have

$$u_0 = \frac{2}{(x-3)^2}$$

$$N(u_k) = \int_0^t (6u_k u_{kx} - u_{kxxx}) dt$$

When k = 0,

$$N(u_0) = u_1 = \int_0^t (6u_0 u_{0x} - u_{0xxx}) dt$$

$$= \int_0^t (-18(x-3)^{-5} + 18(x-3)^{-5}) dt = 0$$

$$u_1 = 0$$

$$u_2 = N(u_0 + u_1) - N(u_0)$$

$$u_2 = \int_0^t (-18(x-3)^{-5} + 18(x-3)^{-5}) dt = 0$$

$$u_3 = 0$$

The successive approximations are:

$$u_0 = \frac{2}{(x-3)^2}$$

$$u_1 = 0$$

$$u_2 = 0$$

$$u_3 = 0$$




Figure 4: The exact solution of example 4.4 (u(x,t)=2/(x-3)<sup>2</sup>)

**CONCLUSION**

In this paper, the analytical solutions of the third-order Korteweg-de Vries (KdV) equations have been obtained using the Variational Iteration Method and the New Iterative Method.

The examples considered showed that Variational Iteration Method and the New Iterative Method are very powerful, efficient and effective methods for solving third order KdV equations because the solutions obtained in series form converge to exact solutions with few iterations.

Hence, solving any other linear and nonlinear differential equations of higher order is very easy by using these powerful methods.

## REFERENCES

- Bahadir A.R. (2005), 'Exponential Finite Difference Method applied to Korteweg-de Vries equation for small times', *Applied Mathematics and Computation*, 160(3), 675-682.
- Batiha B (2009) 'A Variation Iterative Method for Solving the nonlinear Klein-Gordon equation', *Australian Journal of Basic and Applied Sciences*, 3(4): 3876-3890, ISSN 1991-8178.
- Bhalekar S. and Daftardar-Gejji V.(2008), 'New iterative method: application to partial differential equations' *Applied Mathematics and Computation*, 203(2):778–783.
- Dehghan M and Shokri A (2007), 'A Numerical Method for KdV equation using collocation and radial basis functions', *Nonlinear Dynamics*, 50, 111-120
- Eljaily M. H. and Tarig M.E. (2015) 'Homotopy Perturbation Transform Method for solving Korteweg-de Vries (kdv) equation,' *Pure and Applied Mathematics Journal*, 4(6), 264-268.
- Foad Saadi, M. Jalali Azizpour and S.A. Zahedi (2010), 'Analytical Solution of Korteweg-de Vries (KdV) Equation', *International Journal of Physical and Mathematical Sciences*, 4(9)
- Kutluay S, Bahadir A. R and Ozer A. (2000), 'A Small Time Solutions for the Korteweg-de Vries equation', *Applied Mathematics and Computation*, 107, 203-210
- Luwai Wazzan (2016) 'Exact solutions for the family of third order Korteweg de-Vries equations', *Communications in Numerical Analysis* 2, 108-117
- Maryam Sarboland and Azim Amnataei (2015), 'On the Numerical Solution of the Nonlinear Korteweg-de Vries Equation', *Systems Science & Control Engineering: An Open Access Journal*, 3, 69-80
- Muhammad Abbas, Muhammad Kashif Iqbal, Bushra Zafar and Shazalina Binti Mat Zin (2019), 'New Cubic B-spline Approximations for solving Non-linear Third-order Korteweg-de Vries Equation' *Indian Journal of Science and Technology*, 12(6) ISSN 0974-6846
- Olufemi Ige, Matti Heilio, Razaq Adekola Oderinu and Tarig M. Elzaki (2019), 'Adomian Polynomial and Elzaki Transform Method of Solving Third Order Korteweg-de Vries equations', *Global Journal of Pure and Applied Mathematics* ISSN 0973-1768 15(3) 261-277
- Olusola Kolebaje and Oluwole Oyewande (2012), 'Numerical Solution of the Korteweg De Vries Equation by Finite Difference and Adomian Decomposition Method', *International Journal of Basic and Applied Sciences*, 1 (3) (2012) 321-33.
- Schiesser W. (1994), 'Method of lines solution of the Korteweg-de Vries equation', *Computers and Mathematics with Applications*, 28(10) 147-154.
- Shraddha S. Chavan and Mihir M. Panchal (2014), 'Solution of Third Order Korteweg-De Vries Equation by Homotopy Perturbation Method Using Elzaki Transform', *International Journal for Research in Applied Science and Engineering Technology (ijraset)*, ISSN: 2321-965. 2(7)
- Siraj U.I, Khattak A.J and Tirmizi I.A (2008), 'A Meshfree Method for Numerical Solution of KdV equation', *Engineering Analysis with Boundary Elements*, 32, 849-855
- Vincent Daniel David, Mojtaba Nazari, Faisal Salah and Zainal Abdul Aziz (2013), 'Approximate Analytical Solution for the Forced Korteweg-de Vries Equation', *Hindawi Publishing Corporation Journal of Applied Mathematics* Article ID 795818, 9 pages
- Wazwaz Abdul-Majid (2009): 'Partial Differential Equations and Solitary Waves Theory' *Higher Education Press, Beijing and Springer-Verlag Berlin Heidelberg*.
- Xiao M.L., Wang R., and Zhu C.H (2011): 'Applying Multiquadric quasi-interpolation to solve KdV equation', *Mathematical Research Exposition*, 31, 191-201